

## ATTACHMENT 2

### QUALITY MEASURES

Measure Category	Measure	Brief Description	Target Performance Percentage
<b>Asthma</b>			
Asthma Care	Use of appropriate Medications for People with Asthma – Total	Use of appropriate medications for people with persistent asthma <b>Note:</b> Exclusions to this measure shall include any member defined in the denominator who has a diagnosis of one of the following: 471 – Nasal polyps 471.0 – Polyp of nasal cavity 471.1 – Polypoid sinus degeneration 471.8 – Other polyp of sinus 471.9 – Unspecified nasal polyp	
<b>Back Pain</b>			
Appropriate Use of Imaging	Use of Imaging Studies for Low Back Pain	The percentage of patients with a primary diagnosis of low back pain that did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of the diagnosis.	
<b>Depression</b>			
Depression	Antidepressant Medication Management - Effective Continuation Phase Treatment	The percentage of patients 18 years of age and older who were diagnosed with a new episode of major depression, treated with antidepressant medication, and who remained on an antidepressant medication treatment.	
<b>Diabetes</b>			
Diabetes	Retinal Eye Exam Performed	Diabetes Care – Eye Exam as measured by percentage of diabetic (type 1 and type 2) members 18-75 who had a comprehensive eye exam during measurement year	
Diabetes	Blood Sugar Screening	HbA1C testing	
Diabetes	Cholesterol Screening	Diabetes Care – Cholesterol Screening as measured by the percentage of diabetic enrollees 18-75 with diabetes (type 1 and type 2) who had an LDL-C screening test performed during the measurement year	Retired; being replaced
Diabetes	Medical Attention for Nephropathy	Diabetes Care – Kidney Disease Monitoring as measured by the percentage of diabetic (type 1 or type 2) members 18-75 who had a kidney function test during the measurement year	
<b>Heart Disease</b>			
Heart Disease	Persistence of Beta-Blocker Treatment after a	Persistent beta-blocker treatment after acute myocardial infarction (AMI) discharge.	

	Heart Attack		
Heart Disease	Cholesterol Screening for Patients with Cardiovascular Conditions	Cardiovascular Care – Cholesterol Screening as measured by the percentage of members 18–75 years of age who were discharged alive for Acute Myocardial Infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) from January 1–November 1 of the year prior to the measurement year, or who had a diagnosis of ischemic vascular disease (IVD) during the measurement year and the year prior to the measurement year, who had an LDL-C screening test performed during the measurement year	Retired; being replaced
<b>Adult BMI</b>			
BMI	Adult	Adult BMI Assessment	
<b>Medication Monitoring</b>			
Medication Monitoring	Annual Monitoring For Patients On Persistent Medications - Total	Therapeutic monitoring for adults on persistent medications	
<b>Respiratory Infections</b>			
Respiratory Infections	Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	Antibiotics were <u>not</u> prescribed for patients with viral bronchitis	
<b>Other Additional Measures</b>			
Member Survey	Member Satisfaction	<b>Member Survey Score:</b> The Member Survey component will consist of a ranking system ranging from one to ten. The Provider will earn zero points for any Member Survey question where the Provider's performance level on a Member Survey question is below eight. The Provider will earn one point for any Member Survey question where the Provider's performance level is eight or greater. The Provider will earn a maximum of one (1) point for any question within the Member Survey. Provider's total Member Survey Score is calculated by dividing the Provider's total number of points for all Member Surveys completed during an Accounting Period (numerator) by the maximum number of eligible points for all Member Surveys completed during	

		an Accounting Period (denominator).	
<b>Women's Health</b>			
Women's Health	Breast Cancer Screening	Adult mammography for breast cancer screening as measured by the percentage of women 40 to 69 years old who had a mammography	
Women's Health	Cervical Cancer Screening	Adult pap smears for cervical cancer screening	
Women's Health	Chlamydia Screening - Total	Chlamydia screening in sexually active females	